

On the k-Fibonacci words

José L. RAMÍREZ

Instituto de Matemáticas y sus
Aplicaciones
Universidad Sergio Arboleda, Colombia
email:
josel.ramirez@ima.usergioarboleda.edu.co

Gustavo N. RUBIANO

Departamento de Matemáticas
Universidad Nacional de Colombia,
Colombia
email: gnrubiano@unal.edu.co

Abstract. In this paper we define the k-Fibonacci words in analogy with the definition of the k-Fibonacci numbers. We study their properties and we associate to this family of words a family of curves with interesting patterns.

1 Introduction

Fibonacci numbers and their generalizations have many interesting properties and applications to almost every fields of science and arts (e.g. see [13]). The Fibonacci numbers F_n are the terms of the sequence $0, 1, 1, 2, 3, 5, \dots$ wherein each term is the sum of the two previous terms, beginning with the values $F_0 = 0$, and $F_1 = 1$.

Besides the usual Fibonacci numbers many kinds of generalizations of these numbers have been presented in the literature. In particular, a generalization is the k-Fibonacci numbers [11].

For any positive real number k , the k-Fibonacci sequence, say $\{F_{k,n}\}_{n \in \mathbb{N}}$ is defined recurrently by

$$F_{k,0} = 0, F_{k,1} = 1 \quad \text{and} \quad F_{k,n+1} = kF_{k,n} + F_{k,n-1}, \quad n \geq 1. \quad (1)$$

Computing Classification System 1998: I.3.7, G.2.0

Mathematics Subject Classification 2010: 11B39, 05A05, 68R15

Key words and phrases: Fibonacci word, k-Fibonacci numbers, k-Fibonacci words, k-Fibonacci curves

In [11], k -Fibonacci numbers were found by studying the recursive application of two geometrical transformations used in the four-triangle longest-edge (4TLE) partition. These numbers have been studied in several papers, see [5, 10, 11, 12, 18, 19, 23].

The characteristic equation associated to the recurrence relation (1) is $x^2 = kx + 1$. The roots of this equation are

$$r_{k,1} = \frac{k + \sqrt{k^2 + 4}}{2}, \quad \text{and} \quad r_{k,2} = \frac{k - \sqrt{k^2 + 4}}{2}.$$

Some of the properties that the k -Fibonacci numbers verify are (see [11, 12] for the proofs).

- Binet Formula: $F_{k,n} = \frac{r_{k,1}^n - r_{k,2}^n}{r_{k,1} - r_{k,2}}$.
- Combinatorial Formula: $F_{k,n} = \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-1-i}{i} k^{n-1-2i}$.
- $\lim_{n \rightarrow \infty} \frac{F_{k,n}}{F_{k,n-1}} = r_{k,1}$.

On the other hand, there is a word-combinatorial interpretation of the Fibonacci sequence. Fibonacci words are words over $\{0, 1\}$ defined recursively as follows:

$$f_0 = 1, \quad f_1 = 0, \quad f_n = f_{n-1}f_{n-2}, \quad n \geq 2.$$

The words f_n are referred to as the finite Fibonacci words and it is clear that $|f_n| = F_{n+1}$. The limit

$$\mathbf{f} = \lim_{n \rightarrow \infty} f_n = 0100101001001010010100100101 \dots$$

is called the Fibonacci word. This word is certainly one of the most studied words in the combinatorics on words, (see, e.g., [2, 6, 7, 9, 15, 22]). It is the archetype of a Sturmian word [14]. This word can be associated with a curve, which has fractal properties obtained from combinatorial properties of $\mathbf{f}$ [3, 16, 21].

In this paper we introduce a family of words $\mathbf{f}_k$ that generalize the Fibonacci word. Specifically, the k -Fibonacci words are words over $\{0, 1\}$ defined inductively as follows

$$f_{k,0} = 0, \quad f_{k,1} = 0^{k-1}1, \quad f_{k,n} = f_{k,n-1}^k f_{k,n-2},$$

for all $n \geq 2$ and $k \geq 1$. Then it is clear that $|f_{k,n}| = F_{k,n+1}$. The infinite word

$$f_k := \lim_{n \rightarrow \infty} f_{k,n}$$

is called the k -Fibonacci word. In connection with this definition, we investigate some new combinatorial properties and we associate a family of curves with interesting patterns.

2 Definitions and notation

The terminology and notations are mainly those of Lothaire [14] and Allouche and Shallit [1].

Let Σ be a finite alphabet, whose elements are called symbols. A word over Σ is a finite sequence of symbols from Σ . The set of all words over Σ , i.e., the free monoid generated by Σ , is denoted by Σ^* . The identity element ϵ of Σ^* is called the empty word and $\Sigma^+ = \Sigma^* \setminus \{\epsilon\}$. For any word $w \in \Sigma^*$, $|w|$ denotes its length, i.e., the number of symbols occurring in w . The length of ϵ is taken to be equal to 0. If $a \in \Sigma$ and $w \in \Sigma^*$, then $|w|_a$ denotes the number of occurrences of a in w .

For two words $u = a_1 a_2 \cdots a_k$ and $v = b_1 b_2 \cdots b_s$ in Σ^* we denote by uv the concatenation of the two words, that is, $uv = a_1 a_2 \cdots a_k b_1 b_2 \cdots b_s$. If $v = \epsilon$ then $u\epsilon = \epsilon u = u$, moreover, by u^n we denote the word $uu \cdots u$ (n times). A word v is a subword (or factor) of u if there exist $x, y \in \Sigma^*$ such that $u = xvy$. If $x = \epsilon$ ($y = \epsilon$), then v is called prefix (suffix) of u .

The reversal of a word $u = a_1 a_2 \cdots a_n$ is the word $u^R = a_n \cdots a_2 a_1$ and $\epsilon^R = \epsilon$. A word u is a palindrome if $u^R = u$.

An infinite word over Σ is a map $u : \mathbb{N} \rightarrow \Sigma$. It is written $u = a_1 a_2 a_3 \cdots$. The set of all infinite words over Σ is denoted by Σ^ω .

Example 1 The word $p = (p_n)_{n \geq 1} = 0110101000101 \cdots$, where $p_n = 1$ if n is a prime number and $p_n = 0$ otherwise, is an example of an infinite word. p is called the characteristic sequence of the prime numbers.

Definition 2 Let Σ and Δ be alphabets. A morphism is a map $h : \Sigma^* \rightarrow \Delta^*$ such that $h(xy) = h(x)h(y)$ for all $x, y \in \Sigma^*$. It is clear that $h(\epsilon) = \epsilon$.

There is a special class of words, with many remarkable properties, the so-called Sturmian words. These words admit several equivalent definitions (see, e.g. [1] or [14]).

Definition 3 Let $\mathbf{w} \in \Sigma^\omega$. We define $P(\mathbf{w}, n)$, the complexity function of $\mathbf{w}$, to be the map that counts, for all integer $n \geq 0$, the number of subwords of length n in $\mathbf{w}$. An infinite word $\mathbf{w}$ is a Sturmian word if $P(\mathbf{w}, n) = n + 1$ for all integer $n \geq 0$.

Since for any Sturmian word $P(\mathbf{w}, 1) = 2$, then Sturmian words are over two symbols. The word $\mathbf{p}$, in Example 1, is not a Sturmian word because $P(\mathbf{p}, 2) = 4$.

Given two real numbers $\alpha, \beta \in \mathbb{R}$ with α irrational and $0 < \alpha < 1, 0 \leq \beta < 1$, we define the infinite word $\mathbf{w} = w_1 w_2 w_3 \cdots$ as

$$w_n = \lfloor (n+1)\alpha + \beta \rfloor - \lfloor n\alpha + \beta \rfloor.$$

The numbers α and β are the slope and the intercept, respectively. This word is called mechanical. The mechanical words are equivalent to Sturmian words [14]. As special case, when $\beta = 0$, we obtain the characteristic words.

Definition 4 Let α be an irrational number with $0 < \alpha < 1$. For $n \geq 1$, define

$$w_\alpha(n) := \lfloor (n+1)\alpha \rfloor - \lfloor n\alpha \rfloor,$$

and

$$\mathbf{w}(\alpha) := w_\alpha(1)w_\alpha(2)w_\alpha(3)\cdots,$$

then $\mathbf{w}(\alpha)$ is called a characteristic word with slope α .

On the other hand, note that every irrational $\alpha \in (0, 1)$ has a unique continued fraction expansion

$$\alpha = [0, a_1, a_2, a_3, \dots] = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}},$$

where each a_i is a positive integer. Let $\alpha = [0, 1 + d_1, d_2, \dots]$ be an irrational number with $d_1 \geq 0$ and $d_n > 0$ for $n > 1$. To the directive sequence $(d_1, d_2, \dots, d_n, \dots)$, we associate a sequence $(s_n)_{n \geq -1}$ of words defined by

$$s_{-1} = 1, \quad s_0 = 0, \quad s_n = s_{n-1}^{d_n} s_{n-2}, \quad (n \geq 1).$$

Such a sequence of words is called a standard sequence. This sequence is related to characteristic words in the following way. Observe that, for any $n \geq 0$, s_n is a prefix of s_{n+1} , which gives meaning to $\lim_{n \rightarrow \infty} s_n$ as an infinite word. In fact, one can prove [14] that each s_n is a prefix of $\mathbf{w}(\alpha)$ for all $n \geq 0$ and

$$\mathbf{w}(\alpha) = \lim_{n \rightarrow \infty} s_n. \quad (2)$$

Example 5 *The infinite Fibonacci word $\mathbf{f} = 0100101001001010 \cdots$ is a Sturmian word [14], exactly $\mathbf{f} = \mathbf{w}\left(\frac{1}{\phi^2}\right)$ where $\phi = \frac{1+\sqrt{5}}{2}$ is the golden ratio.*

Definition 6 *The Fibonacci morphism $\sigma : \{0, 1\} \rightarrow \{0, 1\}$ is defined by $\sigma(0) = 01$ and $\sigma(1) = 0$.*

The Fibonacci word $\mathbf{f}$ satisfies that $\lim_{n \rightarrow \infty} \sigma^n(1) = \mathbf{f}$ [1].

3 The k -Fibonacci words

Definition 7 *The n th k -Fibonacci words are words over $\{0, 1\}$ defined inductively as follows*

$$f_{k,0} = 0, \quad f_{k,1} = 0^{k-1}1, \quad f_{k,n} = f_{k,n-1}^k f_{k,n-2},$$

for all $n \geq 2$ and $k \geq 1$. The infinite word

$$\mathbf{f}_k := \lim_{n \rightarrow \infty} f_{k,n}$$

is called the k -Fibonacci word.

It is clear that $|f_{k,n}| = F_{k,n+1}$. For $k = 1$ we have the word $\bar{\mathbf{f}} = 1011010110110 \dots$, where $\bar{\mathbf{a}}$ is a morphism, with $\mathbf{a} \in \{0, 1\}$, defined by $\bar{0} = 1, \bar{1} = 0$.

Example 8 *The first k -Fibonacci words are*

$$\begin{aligned} f_1 &= 1011010110110 \cdots = \bar{\mathbf{f}}, & f_2 &= 0101001010010 \cdots, & f_3 &= 0010010010001 \cdots, \\ f_4 &= 0001000100010 \cdots, & f_5 &= 0000100001000 \cdots, & f_6 &= 0000010000010 \cdots. \end{aligned}$$

Definition 9 *The k -Fibonacci morphism $\sigma_k : \{0, 1\} \rightarrow \{0, 1\}$ is defined by $\sigma_k(0) = 0^{k-1}1$ and $\sigma_k(1) = 0^{k-1}10$.*

Theorem 10 *For all $n \geq 0$, $\sigma_k^n(0) = f_{k,n}$ and $\sigma_k^{n+1}(1) = f_{k,n+1}f_{k,n}$. Hence, the k -Fibonacci word $\mathbf{f}_k$ satisfies that $\lim_{n \rightarrow \infty} \sigma_k^n(0) = \mathbf{f}_k$.*

Proof. We prove the two assertions about σ_k^n by induction on n . They are clearly true for $n = 0, 1$. Assume for all $j < n$; we prove them for n :

$$\begin{aligned}\sigma_k^{n+1}(0) &= \sigma_k^n(0^{k-1}1) = (\sigma_k^n(0))^{k-1}\sigma_k^n(1) = f_{k,n}^{k-1}f_{k,n}f_{k,n-1} = f_{k,n}^k f_{k,n-1} = f_{k,n+1}. \\ \sigma_k^{n+2}(1) &= \sigma_k^{n+1}(0^{k-1}10) = (\sigma_k^{n+1}(0))^{k-1}\sigma_k^{n+1}(1)\sigma_k^{n+1}(0) = f_{k,n+1}^{k-1}f_{k,n+1}f_{k,n}f_{k,n+1} \\ &= f_{k,n+1}^k f_{k,n}f_{k,n+1} = f_{k,n+2}f_{k,n+1}. \quad \square\end{aligned}$$

Proposition 11

1. $|f_{k,n}|_1 = F_{k,n}$ and $|f_{k,n+1}|_0 = F_{k,n+1} + F_{k,n}$ for all $n \geq 0$.
2. $\lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} = \frac{r_{k,1}^2}{1 + r_{k,1}}$.
3. $\lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} = r_{k,1}$.
4. $\lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} = 1 + \frac{1}{r_{k,1}}$.

Proof.

1. It is clear by induction on n .

$$\begin{aligned}2. \quad \lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} &= \lim_{n \rightarrow \infty} \frac{F_{k,n+1}}{F_{k,n} + F_{k,n-1}} = \lim_{n \rightarrow \infty} \frac{\frac{F_{k,n+1}}{F_{k,n}}}{1 + \frac{F_{k,n-1}}{F_{k,n}}} = \frac{r_{k,1}^2}{1 + r_{k,1}}. \\ 3. \quad \lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} &= \lim_{n \rightarrow \infty} \frac{F_{k,n+1}}{F_{k,n}} = r_{k,1}. \\ 4. \quad \lim_{n \rightarrow \infty} \frac{|f_{k,n}|_0}{|f_{k,n}|_1} &= \lim_{n \rightarrow \infty} \frac{F_{k,n} + F_{k,n-1}}{F_{k,n}} = 1 + \frac{1}{r_{k,1}}. \quad \square\end{aligned}$$

Proposition 12 *The k -Fibonacci word and the n th k -Fibonacci word satisfy that*

1. *The word 11 is not a subword of the k -Fibonacci word, $k \geq 2$.*
2. *Let ab be the last two symbols of $f_{k,n}$. For $n \geq 1$, we have $ab = 10$ if n is even and $ab = 01$ if n is odd, $k \geq 2$.*
3. *The concatenation of two successive k -Fibonacci words is “almost commutative”, i.e., $f_{k,n-1}f_{k,n-2}$ and $f_{k,n-2}f_{k,n-1}$ have a common prefix the length $F_{k,n} + F_{k,n-1} - 2$ for all $n \geq 2$.*

Proof.

1. It suffices to prove that 11 is not a subword of $f_{k,n}$, for all $n \geq 0$. By induction on n . For $n = 0, 1$ it is clear. Assume for all $j < n$; we prove it for n . We know that $f_{k,n} = f_{k,n-1}^k f_{k,n-2}$ so by the induction hypothesis we have that 11 is not a subword of $f_{k,n-1}$ and $f_{k,n-2}$. Therefore, the only possibility is that 1 is a suffix and prefix of $f_{k,n-1}$ or 1 is a suffix of $f_{k,n-1}$ and a prefix of $f_{k,n-2}$, both there are impossible.
2. By induction on n . For $n = 1, 2$ it is clear. Assume for all $j < n$; we prove it for n . We know that $f_{k,n+1} = f_{k,n}^k f_{k,n-1}$, if $n+1$ is even then by the induction hypothesis the last two symbols of $f_{k,n-1}$ are 10, therefore the last two symbols of $f_{k,n+1}$ are 10. Analogously, if $n+1$ is odd.
3. By induction on n . For $n = 1, 2$ it is clear. Assume for all $j < n$; we prove it for n . By definition of $f_{k,n}$, we have

$$\begin{aligned} f_{k,n-1} f_{k,n-2} &= f_{k,n-2}^k f_{k,n-3} \cdot f_{k,n-3}^k f_{k,n-4} \\ &= (f_{k,n-3}^k f_{k,n-4})^k \cdot f_{k,n-3}^k f_{k,n-3} f_{k,n-4}, \end{aligned}$$

and

$$\begin{aligned} f_{k,n-2} f_{k,n-1} &= f_{k,n-3}^k f_{k,n-4} \cdot f_{k,n-2}^k f_{k,n-3} \\ &= f_{k,n-3}^k f_{k,n-4} \cdot (f_{k,n-3}^k f_{k,n-4})^k \cdot f_{k,n-3} \\ &= (f_{k,n-3}^k f_{k,n-4})^k f_{k,n-3}^k f_{k,n-4} f_{k,n-3}. \end{aligned}$$

Hence the words have a common prefix of length $k(kF_{k,n-2} + F_{k,n-3}) + kF_{k,n-2} = k(F_{k,n-1} + F_{k,n-2})$. By the induction hypothesis $f_{k,n-3} f_{k,n-4}$ and $f_{k,n-4} f_{k,n-3}$ have a common prefix of length $F_{k,n-2} + F_{k,n-3} - 2$. Therefore the words have a common prefix of length

$$k(F_{k,n-1} + F_{k,n-2}) + F_{k,n-2} + F_{k,n-3} - 2 = F_{k,n} + F_{k,n-1} - 2. \quad \square$$

Definition 13 Let $\Phi : \{0, 1\}^* \rightarrow \{0, 1\}^*$ be a map such that Φ deletes the last two symbols, i.e., $\Phi(a_1 a_2 \cdots a_n) = a_1 a_2 \cdots a_{n-2}$ ($n \geq 2$).

Corollary 14 The n th k -Fibonacci word, satisfy for all $n \geq 2$ that

1. $\Phi(f_{k,n-1} f_{k,n-2}) = \Phi(f_{k,n-2} f_{k,n-1})$.
2. $\Phi(f_{k,n-1} f_{k,n-2}) = f_{k,n-2} \Phi(f_{k,n-1}) = f_{k,n-1} \Phi(f_{k,n-2})$.
3. If $f_{k,n} = \Phi(f_{k,n}) ab$, then $\Phi(f_{k,n-2}) ab \Phi(f_{k,n-1}) = f_{k,n-1} \Phi(f_{k,n-2})$.

4. If $f_{k,n} = \Phi(f_{k,n})ab$, then $\Phi(f_{k,n-2})(ab\Phi(f_{k,n-1}))^k = \Phi(f_{k,n})$.

Proof. Parts (a) and (b) follow immediately from Proposition 12-(3) and because of $|f_{k,n}| \geq 2$ for all $n \geq 2$. (c) In fact, if $f_{k,n} = \Phi(f_{k,n})ab$ then from Proposition 12-(2) we have $f_{k,n-2} = \Phi(f_{k,n-2})ab$. Hence $\Phi(f_{k,n-2})ab\Phi(f_{k,n-1}) = f_{k,n-2}\Phi(f_{k,n-1}) = f_{k,n-1}\Phi(f_{k,n-2})$. (d) It is clear from (c) and definition of $f_{k,n}$. $\square$

Theorem 15 $\Phi(f_{k,n})$ is a palindrome for all $n \geq 1$ and $k \geq 1$.

Proof. By induction on n . If $n = 2$ then $\Phi(f_{k,2}) = (0^{k-1}1)^{k-1}0^{k-1}$ is a palindrome. Now suppose that the result is true for all $j < n$; we prove it for n .

$$\begin{aligned} (\Phi(f_{k,n}))^R &= (\Phi(f_{k,n-1}^k f_{k,n-2}))^R = (f_{k,n-1}^k \Phi(f_{k,n-2}))^R = \Phi(f_{k,n-2})^R (f_{k,n-1}^k)^R \\ &= \Phi(f_{k,n-2})(f_{k,n-1}^R)^k. \end{aligned}$$

If n is even then $f_{k,n} = \Phi(f_{k,n})10$ and from Corollary 14-(4), we have that

$$\begin{aligned} (\Phi(f_{k,n}))^R &= \Phi(f_{k,n-2})((\Phi(f_{k,n-1})01)^R)^k = \Phi(f_{k,n-2})(10(\Phi(f_{k,n-1}))^R)^k \\ &= \Phi(f_{k,n-2})(10\Phi(f_{k,n-1}))^k = \Phi(f_{k,n}). \end{aligned}$$

If n is odd, the proof is analogous. $\square$

Corollary 16 1. If $f_{k,n} = \Phi(f_{k,n})ab$ then $ba\Phi(f_{k,n})ab$ is a palindrome.
2. If u is a subword of the k -Fibonacci word, then so is its reversal, u^R .

Theorem 17 Let $\alpha = [0, \bar{k}]$ be an irrational number, with k a positive integer, then

$$w(\alpha) = f_k.$$

Proof. Let $\alpha = [0, \bar{k}]$ an irrational number, then its associated standard sequence is

$$s_{-1} = 1, \quad s_0 = 0, \quad s_1 = s_0^{k-1} s_{-1} = 0^{k-1} 1 \text{ and } s_n = s_{n-1}^k s_{n-2}, \quad n \geq 2.$$

Hence $\{s_n\}_{n \geq 0} = \{f_{k,n}\}_{n \geq 0}$ and from equation (2), we have

$$w(\alpha) = \lim_{n \rightarrow \infty} s_n = f_k. \quad \square$$

Remark. Note that

$$[0, \overline{k}] = \frac{1}{k + \frac{1}{k + \frac{1}{k + \frac{1}{\ddots}}}} = \frac{-k + \sqrt{k^2 + 4}}{2} = -r_{k,2}$$

From the above theorem, we conclude that k -Fibonacci words are Sturmian words.

A fractional power is a word of the form $z = x^n y$, where $n \in \mathbb{Z}^+$ and $x \in \Sigma^+$, and y is power prefix of x . If $|z| = p$ and $|x| = q$, we say that z is a p/q -power, or $z = x^{p/q}$. In the expression $x^{p/q}$, the number p/q is the power's exponent. For example, 01201201 is a $8/3$ -power, $01201201 = (012)^{8/3}$. The index of an infinite word $w \in \Sigma^\omega$ is defined by

$$\text{Ind}(w) := \sup\{r \in \mathbb{Q}_{\geq 1} : w \text{ contains an } r\text{-power}\}$$

For example $\text{Ind}(\mathbf{f}) > 3$ because the cube $(010)^3$ occurs in $\mathbf{f}$ at position 6. In [15] the authors proof that $\text{Ind}(\mathbf{f}) = 2 + \phi \approx 3.618$. A general formula for the index of a Sturmian word was given in [8].

Theorem 18 *If $\mathbf{u}$ is a Sturmian word of slope $\alpha = [0, a_1, a_2, a_3, \dots]$, then*

$$\text{Ind}(w) = \sup_{n \geq 0} \left\{ 2 + a_{n+1} + \frac{q_{n-1} - 2}{q_n} \right\},$$

where q_n is the denominator of $\alpha = [0, a_1, a_2, a_3, \dots, a_n]$ and satisfies $q_{-1} = 0, q_0 = 1, q_{n+1} = a_{n+1}q_n + q_{n-1}$.

Corollary 19 *The index of k -Fibonacci words is $\text{Ind}(\mathbf{f}_k) = 2 + k + \frac{1}{r_{k,1}}$.*

Proof. $\mathbf{f}_k$ is a Sturmian word of slope $\alpha = [0, \overline{k}]$, then it is clear that $q_n = F_{k,n+1}$, and from above theorem

$$\text{Ind}(\mathbf{f}_k) = \sup_{n \geq 0} \left\{ 2 + k + \frac{F_{k,n} - 2}{F_{k,n+1}} \right\} = 2 + k + \frac{1}{r_{k,1}}. \quad \square$$

4 The k-Fibonacci Word Curve

The Fibonacci word can be associated to a curve from a drawing rule. We must travel the word in a particular way, depending on the symbol read a particular action is produced, this idea is the same as that used in the L-Systems [17]. In this case, the drawing rule is called “odd-even drawing rule” [16], this is defined as shown in the following table.

Symbol	Action
1	Draw a line forward.
0	Draw a line forward and if the symbol 0 is in a position even then turn θ degree and if 0 is in a position odd then turn $-\theta$ degrees.

Definition 20 The n th-curve of Fibonacci, denoted by $\mathcal{F}_n$, is obtained by applying the odd-even drawing rule to the word f_n . The Fibonacci word fractal $\mathcal{F}$, is defined as

$$\mathcal{F} := \lim_{n \rightarrow \infty} \mathcal{F}_n.$$

Example 21 In Figure 1 we show the curve $\mathcal{F}_{10}$ and $\mathcal{F}_{17}$. The graphics in this paper were generated using the software **Mathematica 9.0**, [20].





Figure 1: Fibonacci curves $\mathcal{F}_{10}$ and $\mathcal{F}_{17}$ corresponding to the words f_{10} and f_{17}

Properties of Fibonacci Word Fractal can be found in [3, 4, 16].

Definition 22 The n th k -curve of Fibonacci, denoted by $\mathcal{F}_{k,n}$, is obtained by applying the odd-even drawing rule to the word $f_{k,n}$. The k -Fibonacci word curve $\mathcal{F}_k$ is defined as

$$\mathcal{F}_k := \lim_{n \rightarrow \infty} \mathcal{F}_{k,n}.$$

Table 1: Some curves $\mathcal{F}_{k,n}$ with $\theta = 90^\circ$

In Table 1, we show some curves $\mathcal{F}_{k,n}$ with an angle $\theta = 90^\circ$.


In Table 2, we show some curves $\mathcal{F}_{k,n}$ with an angle $\theta = 60^\circ$. In general these curves have a lot of patterns because the index is large, see Corollary 19.

Proposition 23 *The k -Fibonacci word curve and the curve $\mathcal{F}_{k,n}$ have the following properties:*

1. The k -Fibonacci curve $\mathcal{F}_k$ is composed only of segments of length 1 or 2.
2. The $\mathcal{F}_{k,n}$ is symmetric.
3. The number of turns in the curve $\mathcal{F}_{k,n}$ is $F_{k,n} + F_{k,n-1}$.
4. If n is even then the $\mathcal{F}_{k,n}$ curve is similar to the curve $\mathcal{F}_{k,n-2}$ and if n is odd then the $\mathcal{F}_{k,n}$ curve is similar to the curve $\mathcal{F}_{k,n-3}$.

Proof.

1. It is clear from Proposition 12-1, because 110 and 111 are not subwords of $\mathbf{f}_k$.
2. It is clear from Theorem 15, because $\mathbf{f}_{k,n} = \Phi(\mathbf{f}_{k,n})\mathbf{ab}$, where $\Phi(\mathbf{f}_{k,n})$ is a palindrome.
3. It is clear from definition of odd-even drawn rule and because $|\mathbf{f}_{k,n+1}|_0 = F_{k,n+1} + F_{k,n}$.

Table 2: Some curves $\mathcal{F}_{k,n}$ with $\theta = 60^\circ$

4. If n is even. It is clear that $\sigma_k^2(f_{k,n-2}) = f_{k,n}$. We are going to prove that σ_k^2 guaranties the odd-even alternation required by the odd-even drawing rule. In fact, $\sigma_k^2(0) = \sigma_k(0^{k-1}1) = (0^{k-1}1)^k 0$ and $\sigma_k^2(1) = (0^{k-1}1)^k 0^k 1$. As k is even, then $|\sigma_k^2(0)|$ and $|\sigma_k^2(1)|$ are odd. Hence if $|w|$ is even (odd) then $|\sigma_k^2(w)|$ is even (odd). Since σ_k^2 preserves the parity of length then any subword in the k -Fibonacci word preserves the parity of position.

Finally, we have to proof that the resulting angle of a pattern must be preserved or inverted by σ_k^2 . Let $\alpha(w)$ be the function that gives the resulting angle of a word w through the odd-even drawing rule of angle

θ . Note that $a(00) = 0^\circ$, $a(01) = -\theta^\circ$ and $a(10) = \theta^\circ$. Therefore

$$\begin{aligned}
 a(\sigma_k^2(00)) &= a((0^{k-1}1)^k 0 (0^{k-1}1)^k 0) \\
 &= a((0^{k-1}1)^k) a(00^{k-1}) a(1(0^{k-1}1)^{k-1}0) \\
 &= -k\theta^\circ + 0^\circ + k\theta^\circ = 0^\circ \\
 a(\sigma_k^2(01)) &= a((0^{k-1}1)^k 0 (0^{k-1}1)^k 0^k 1) \\
 &= a((0^{k-1}1)^k) a(00^{k-1}) a(1(0^{k-1}1)^{k-1}0) a(0^{k-1}1) \\
 &= -k\theta^\circ + 0^\circ + k\theta^\circ - \theta^\circ = -\theta^\circ \\
 a(\sigma_k^2(10)) &= a((0^{k-1}1)^k 0^k 1 (0^{k-1}1)^k 0) \\
 &= a((0^{k-1}1)^k) a(0^k) a(1(0^{k-1}1)^k 0) \\
 &= -k\theta^\circ + 0^\circ + (k+1)\theta^\circ = \theta^\circ
 \end{aligned}$$

Then σ_k^2 inverts the resulting angle, i.e., $a(w) = -a(\sigma_k^2(w))$ for any word w . Therefore the image of a pattern by σ_k^2 is the rotation of this pattern by a rotation of $-\theta^\circ$. Since $\sigma_k^2(f_{k,n-2}) = f_{k,n}$, then the curve $\mathcal{F}_{k,n}$ is similar to the curve $\mathcal{F}_{k,n-2}$.

If n is odd the proof is similar, but using σ_k^3 . □

Example 24 In Figure 2 $\mathcal{F}_{4,4}$ looks similar to $\mathcal{F}_{4,6}$, $\mathcal{F}_{4,8}$ and so on.




Figure 2: Curves $\mathcal{F}_{4,4}$, $\mathcal{F}_{4,6}$, $\mathcal{F}_{4,8}$ with $\theta = 60^\circ$

In Figure 3 $\mathcal{F}_{5,3}$ looks similar to $\mathcal{F}_{5,6}$.

Acknowledgements

The first author was partially supported by Universidad Sergio Arboleda under grant number USA-II-2012-14.




Figure 3: Curves $\mathcal{F}_{5,3}, \mathcal{F}_{5,6}$ with $\theta = 60^\circ$

References

- [1] J. Allouche, J. Shallit, *Automatic Sequences*, Cambridge University Press, 2003. $\Rightarrow$ 214, 216
- [2] J. Berstel, Fibonacci words—a survey, in: G. Rosenberg, A. Salomaa (Eds.), *The Book of L*, Springer, Berlin, 1986, pp. 11–26. $\Rightarrow$ 213
- [3] A. Blondin-Mass, S. Brlek, A. Garon, S. Labb, Two infinite families of polyominoes that tile the plane by translation in two distinct ways, *Theoret. Comput. Sci.* **412** (2011) 4778–4786. $\Rightarrow$ 213, 221
- [4] A. Blondin-Mass, S. Brlek, S. Labb, M. Mends France, Complexity of the Fibonacci snowflake, *Fractals* **20** (2012) 257–260. $\Rightarrow$ 221
- [5] C. Bolat, H. Kse, On the properties of k -Fibonacci numbers, *Int. J. Contemp. Math. Sciences*, **22**, 5 (2010) 1097–1105. $\Rightarrow$ 213
- [6] J. Cassaigne, On extremal properties of the Fibonacci word, *RAIRO - Theor. Inf. Appl.* **42**, (4) (2008) 701–715. $\Rightarrow$ 213
- [7] W. Chuan, Fibonacci words, *Fibonacci Quart.*, **30**, 1 (1992) 68–76. $\Rightarrow$ 213
- [8] D. Damanik, D. Lenz, The index of Sturmian sequences, *European J. Combin.*, **23** (2002) 23–29. $\Rightarrow$ 220
- [9] A. de Luca, Sturmian words: structure, combinatorics, and their arithmetics, *Theor. Comput. Sci.* **183**, 1 (1997) 45–82. $\Rightarrow$ 213
- [10] S. Falcon, A. Plaza, On k -Fibonacci sequences and polynomials and their derivatives, *Chaos Solitons Fractals* **39**, 3 (2009) 1005–1019. $\Rightarrow$ 213
- [11] S. Falcon, A. Plaza, On the Fibonacci k -numbers, *Chaos Solitons Fractals* **32**, 5 (2007) 1615–1624. $\Rightarrow$ 212, 213
- [12] S. Falcon, A. Plaza, The k -Fibonacci sequence and the Pascal 2-triangle, *Chaos Solitons Fractals* **33**, 1 (2007) 38–49. $\Rightarrow$ 213
- [13] T. Koshy, *Fibonacci and Lucas numbers with Applications*, Wiley-Interscience, 2001. $\Rightarrow$ 212
- [14] M. Lothaire, *Algebraic Combinatorics on Words*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2002. $\Rightarrow$ 213, 214, 215, 216

- [15] F. Mignosi, G. Pirillo, Repetitions in the Fibonacci infinite word, *RAIRO – Theor. Inf. Appl.* **26** (1992) 199–204. $\Rightarrow$ 213, 220
- [16] A. Monnerot, The Fibonacci word fractal, preprint, 2009. $\Rightarrow$ 213, 221
- [17] P. Prusinkiewicz, A. Lindenmayer, *The Algorithmic Beauty of Plants*, Springer-Verlag, Nueva York, 2004. $\Rightarrow$ 221
- [18] J. Ramírez, Incomplete k-Fibonacci and k-Lucas numbers, *Chinese Journal of Mathematics* (2013). $\Rightarrow$ 213
- [19] J. Ramírez. Some properties of convolved k-Fibonacci numbers. *ISRN Combinatorics* (2013) ID759641, 5pp. $\Rightarrow$ 213
- [20] J. Ramírez, G. Rubiano, Generating fractals curves from homomorphisms between languages [with Mathematica[®]] (Spanish), *Rev. Integr. Temas Mat.* **30**, 2 (2012) 129–150. $\Rightarrow$ 221
- [21] J. Ramírez, G. Rubiano, R. de Castro, A generalization of the Fibonacci word fractal and the Fibonacci snowflake, preprint arXiv:1212.1368, 2013. $\Rightarrow$ 213
- [22] W. Rytter, The structure of subword graphs and suffix trees of Fibonacci words, *Theoret. Comput. Sci.* **363**, 2 (2006) 211–223. $\Rightarrow$ 213
- [23] A. Salas. About k-Fibonacci numbers and their associated numbers, *Int. Math. Forum.* **50**, 6 (2011) 2473–2479. $\Rightarrow$ 213

Received: June 12, 2013 • Revised: October 25, 2013