



# Multiplicative inequalities for weighted arithmetic and harmonic operator means

Sever S. Dragomir

Mathematics, College of Engineering & Science  
Victoria University, Melbourne City, Australia

email: sever.dragomir@vu.edu.au  
URL: <http://rgmia.org/dragomir>

School of Computer Science & Applied Mathematics,  
University of the Witwatersrand,  
Johannesburg, South Africa

**Abstract.** In this paper we establish some multiplicative inequalities for weighted arithmetic and harmonic operator means under various assumption for the positive invertible operators  $A$ ,  $B$ . Some applications when  $A$ ,  $B$  are bounded above and below by positive constants are given as well.

## 1 Introduction

Throughout this paper  $A$ ,  $B$  are positive invertible operators on a complex Hilbert space  $(H, \langle \cdot, \cdot \rangle)$ . We use the following notations for operators

$$A\nabla_v B := (1 - v)A + vB,$$

the *weighted operator arithmetic mean*,

$$A\sharp_v B := A^{1/2} \left( A^{-1/2} B A^{-1/2} \right)^v A^{1/2},$$

---

**2010 Mathematics Subject Classification:** 47A63, 47A30, 15A60, 26D15, 26D10

**Key words and phrases:** Young's inequality, convex functions, arithmetic mean-harmonic mean inequality, operator means, operator inequalities

the *weighted operator geometric mean* and

$$A!_v B := \left( (1-v) A^{-1} + v B^{-1} \right)^{-1}$$

the *weighted operator harmonic mean*, where  $v \in [0, 1]$ .

When  $v = \frac{1}{2}$ , we write  $A\nabla B$ ,  $A\sharp B$  and  $A!B$  for brevity, respectively.

The following fundamental inequality between the weighted arithmetic, geometric and harmonic operator means holds

$$A!_v B \leq A\sharp_v B \leq A\nabla_v B \quad (1)$$

for any  $v \in [0, 1]$ .

For various recent inequalities between these means we recommend the recent papers [3]-[6], [8]-[12] and the references therein.

The following additive double inequality has been obtained in the recent paper [7]:

$$v(1-v) \frac{(b-a)^2}{\max\{b,a\}} \leq A_v(a,b) - H_v(a,b) \leq v(1-v) \frac{(b-a)^2}{\min\{b,a\}}, \quad (2)$$

for any  $a, b > 0$  and  $v \in [0, 1]$ , where  $A_v(a,b)$  and  $H_v(a,b)$  are the scalar weighted arithmetic mean and harmonic mean, respectively, namely

$$A_v(a,b) := (1-v)a + vb \text{ and } H_v(a,b) := \frac{ab}{(1-v)b + va}.$$

In particular,

$$\frac{1}{4} \frac{(b-a)^2}{\max\{b,a\}} \leq A(a,b) - H(a,b) \leq \frac{1}{4} \frac{(b-a)^2}{\min\{b,a\}}, \quad (3)$$

where

$$A(a,b) := \frac{a+b}{2} \text{ and } H(a,b) := \frac{2ab}{b+a}.$$

We consider the *Kantorovich's constant* defined by

$$K(h) := \frac{(h+1)^2}{4h}, \quad h > 0. \quad (4)$$

The function  $K$  is decreasing on  $(0, 1)$  and increasing on  $[1, \infty)$ ,  $K(h) \geq 1$  for any  $h > 0$  and  $K(h) = K(\frac{1}{h})$  for any  $h > 0$ .

Observe that for any  $h > 0$

$$K(h) - 1 = \frac{(h-1)^2}{4h} = K\left(\frac{1}{h}\right) - 1.$$

Observe that

$$K\left(\frac{b}{a}\right) - 1 = \frac{(b-a)^2}{4ab} \text{ for } a, b > 0.$$

Since, obviously

$$ab = \min\{a, b\} \max\{a, b\} \text{ for } a, b > 0,$$

then we have the following version of (2):

$$\begin{aligned} 4\nu(1-\nu) \min\{a, b\} \left[ K\left(\frac{b}{a}\right) - 1 \right] &\leq A_\nu(a, b) - H_\nu(a, b) \\ &\leq 4\nu(1-\nu) \max\{a, b\} \left[ K\left(\frac{b}{a}\right) - 1 \right]. \end{aligned} \quad (5)$$

for any  $a, b > 0$  and  $\nu \in [0, 1]$ .

For positive invertible operators  $A, B$  we define

$$A\nabla_\infty B := \frac{1}{2}(A+B) + \frac{1}{2}A^{1/2} \left| A^{-1/2}(B-A)A^{-1/2} \right| A^{1/2}$$

and

$$A\nabla_{-\infty} B := \frac{1}{2}(A+B) - \frac{1}{2}A^{1/2} \left| A^{-1/2}(B-A)A^{-1/2} \right| A^{1/2}.$$

If we consider the continuous functions  $f_\infty, f_{-\infty} : [0, \infty) \rightarrow [0, \infty)$  defined by

$$f_\infty(x) = \max\{x, 1\} = \frac{1}{2}(x+1) + \frac{1}{2}|x-1|$$

and

$$f_{-\infty}(x) = \max\{x, 1\} = \frac{1}{2}(x+1) - \frac{1}{2}|x-1|,$$

then, obviously, we have

$$A\nabla_{\pm\infty} B = A^{1/2} f_{\pm\infty} \left( A^{-1/2} B A^{-1} \right) A^{1/2}.$$

If  $A$  and  $B$  are commutative, then

$$A\nabla_{\pm\infty} B = \frac{1}{2}(A+B) \pm \frac{1}{2}|B-A| = B\nabla_{\pm\infty} A.$$

The following additive inequality between the weighted arithmetic and harmonic operator means holds [7]:

**Theorem 1** Let  $A, B$  be positive invertible operators and  $M > m > 0$  such that the condition

$$mA \leq B \leq MA \quad (6)$$

holds. Then we have

$$\begin{aligned} 4\nu(1-\nu)g(m, M)A\nabla_{-\infty}B &\leq A\nabla_\nu B - A!\nu B \\ &\leq 4\nu(1-\nu)G(m, M)A\nabla_\infty B, \end{aligned} \quad (7)$$

where

$$g(m, M) := \begin{cases} K(M) - 1 & \text{if } M < 1, \\ 0 & \text{if } m \leq 1 \leq M, \\ K(m) - 1 & \text{if } 1 < m \end{cases}$$

and

$$G(m, M) := \begin{cases} K(m) - 1 & \text{if } M < 1, \\ \max\{K(m), K(M)\} - 1 & \text{if } m \leq 1 \leq M, \\ K(M) - 1 & \text{if } 1 < m. \end{cases}$$

In particular,

$$g(m, M)A\nabla_{-\infty}B \leq A\nabla B - A!B \leq G(m, M)A\nabla_\infty B. \quad (8)$$

Motivated by the above facts, we establish in this paper some multiplicative inequalities for weighted arithmetic and harmonic operator means under various assumption for the positive invertible operators  $A, B$ . Some applications when  $A, B$  are bounded above and below by positive constants are given as well.

## 2 Multiplicative inequalities

The following result is of interest in itself:

**Lemma 1** For any  $a, b > 0$  and  $\nu \in [0, 1]$  we have

$$\nu(1-\nu) \left(1 - \frac{\min\{a, b\}}{\max\{a, b\}}\right)^2 \leq \frac{A_\nu(a, b)}{H_\nu(a, b)} - 1 \leq \nu(1-\nu) \left(\frac{\max\{a, b\}}{\min\{a, b\}} - 1\right)^2. \quad (9)$$

In particular,

$$\frac{1}{4} \left(1 - \frac{\min\{a, b\}}{\max\{a, b\}}\right)^2 \leq \frac{A(a, b)}{H(a, b)} - 1 \leq \frac{1}{4} \left(\frac{\max\{a, b\}}{\min\{a, b\}} - 1\right)^2. \quad (10)$$

**Proof.** We have for any  $a, b > 0$  and  $\nu \in [0, 1]$  that

$$\begin{aligned} \frac{A_\nu(a, b)}{H_\nu(a, b)} &= \frac{[(1-\nu)a + \nu b][(1-\nu)b + \nu a]}{ab} \\ &= \frac{(1-\nu)^2 ab + \nu(1-\nu)b^2 + \nu(1-\nu)a^2 + \nu^2 ab}{ab} \\ &= \frac{\nu(1-\nu)(b^2 + a^2) + (1-2\nu+2\nu^2)ab}{ab}, \end{aligned}$$

which is equivalent with

$$\frac{A_\nu(a, b)}{H_\nu(a, b)} - 1 = \nu(1-\nu) \frac{(b-a)^2}{ab} \quad (11)$$

for any  $a, b > 0$  and  $\nu \in [0, 1]$ .

Since  $\min^2\{a, b\} \leq ab \leq \max^2\{a, b\}$  hence

$$\begin{aligned} \nu(1-\nu) \frac{(b-a)^2}{ab} &\leq \nu(1-\nu) \frac{(b-a)^2}{\min^2\{a, b\}} \\ &= \nu(1-\nu) \left( \frac{\max\{a, b\}}{\min\{a, b\}} - 1 \right)^2 \end{aligned}$$

and

$$\begin{aligned} \nu(1-\nu) \frac{(b-a)^2}{ab} &\geq \nu(1-\nu) \frac{(b-a)^2}{\max^2\{a, b\}} \\ &= \nu(1-\nu) \left( 1 - \frac{\min\{a, b\}}{\max\{a, b\}} \right)^2 \end{aligned}$$

and by (11) we get the desired result (9).  $\square$

We observe that the inequality (9) can be written in an equivalent form as

$$\begin{aligned} &\left[ \nu(1-\nu) \left( 1 - \frac{\min\{a, b\}}{\max\{a, b\}} \right)^2 + 1 \right] H_\nu(a, b) \\ &\leq A_\nu(a, b) \\ &\leq \left[ \nu(1-\nu) \left( \frac{\max\{a, b\}}{\min\{a, b\}} - 1 \right)^2 + 1 \right] H_\nu(a, b) \end{aligned} \quad (12)$$

for any  $a, b > 0$  and  $\nu \in [0, 1]$ , while (10) as

$$\begin{aligned} & \left[ \frac{1}{4} \left( 1 - \frac{\min\{a, b\}}{\max\{a, b\}} \right)^2 + 1 \right] H(a, b) \\ & \leq A(a, b) \\ & \leq \left[ \frac{1}{4} \left( \frac{\max\{a, b\}}{\min\{a, b\}} - 1 \right)^2 + 1 \right] H(a, b) \end{aligned} \quad (13)$$

for any  $a, b > 0$ .

**Corollary 1** *For any  $a, b \in [k, K] \subset (0, \infty)$  and  $\nu \in [0, 1]$  we have*

$$\frac{A_\nu(a, b)}{H_\nu(a, b)} - 1 \leq \nu(1 - \nu) \left( \frac{K}{k} - 1 \right)^2. \quad (14)$$

In particular,

$$\frac{A(a, b)}{H(a, b)} - 1 \leq \frac{1}{4} \left( \frac{K}{k} - 1 \right)^2. \quad (15)$$

We have the following multiplicative inequality between the weighted arithmetic and harmonic operator means:

**Theorem 2** *Let  $A, B$  be positive invertible operators and  $M > m > 0$  such that the condition (6) holds. Then we have*

$$\begin{aligned} & \left[ \nu(1 - \nu) \left( 1 - \frac{\min\{M, 1\}}{\max\{m, 1\}} \right)^2 + 1 \right] A!_\nu B \\ & \leq A \nabla_\nu B \\ & \leq \left[ \nu(1 - \nu) \left( \frac{\max\{M, 1\}}{\min\{m, 1\}} - 1 \right)^2 + 1 \right] A!_\nu B \end{aligned} \quad (16)$$

for any  $\nu \in [0, 1]$ .

In particular,

$$\begin{aligned} & \left[ \frac{1}{4} \left( 1 - \frac{\min\{M, 1\}}{\max\{m, 1\}} \right)^2 + 1 \right] A!B \\ & \leq A \nabla B \\ & \leq \left[ \frac{1}{4} \left( \frac{\max\{M, 1\}}{\min\{m, 1\}} - 1 \right)^2 + 1 \right] A!B. \end{aligned} \quad (17)$$

**Proof.** If we write the inequality (12) for  $a = 1$  and  $b = x \in (0, \infty)$  then we have

$$\begin{aligned} & \left[ v(1-v) \left( 1 - \frac{\min\{1, x\}}{\max\{1, x\}} \right)^2 + 1 \right] (1-v+vx^{-1})^{-1} \\ & \leq 1-v+vx \\ & \leq \left[ v(1-v) \left( \frac{\max\{1, x\}}{\min\{1, x\}} - 1 \right)^2 + 1 \right] (1-v+vx^{-1})^{-1}. \end{aligned} \quad (18)$$

for any  $v \in [0, 1]$ .

If  $x \in [m, M] \subset (0, \infty)$ , then  $\max\{m, 1\} \leq \max\{x, 1\} \leq \max\{M, 1\}$  and  $\min\{m, 1\} \leq \min\{x, 1\} \leq \min\{M, 1\}$ .

We have

$$\left( \frac{\max\{1, x\}}{\min\{1, x\}} - 1 \right)^2 \leq \left( \frac{\max\{M, 1\}}{\min\{m, 1\}} - 1 \right)^2$$

and

$$\left( 1 - \frac{\min\{M, 1\}}{\max\{m, 1\}} \right)^2 \leq \left( 1 - \frac{\min\{1, x\}}{\max\{1, x\}} \right)^2$$

for any  $x \in [m, M] \subset (0, \infty)$ .

Therefore, by (18) we have

$$\begin{aligned} & \left[ v(1-v) \left( 1 - \frac{\min\{M, 1\}}{\max\{m, 1\}} \right)^2 + 1 \right] (1-v+vx^{-1})^{-1} \\ & \leq 1-v+vx \\ & \leq \left[ v(1-v) \left( \frac{\max\{M, 1\}}{\min\{m, 1\}} - 1 \right)^2 + 1 \right] (1-v+vx^{-1})^{-1}, \end{aligned} \quad (19)$$

for any  $x \in [m, M]$  and for any  $v \in [0, 1]$ .

If we use the continuous functional calculus for the positive invertible operator  $X$  with  $mI \leq X \leq MI$ , then we have from (19) that

$$\begin{aligned} & \left[ v(1-v) \left( 1 - \frac{\min\{M, 1\}}{\max\{m, 1\}} \right)^2 + 1 \right] ((1-v)I+vx^{-1})^{-1} \\ & \leq (1-v)I+vx \\ & \leq \left[ v(1-v) \left( \frac{\max\{M, 1\}}{\min\{m, 1\}} - 1 \right)^2 + 1 \right] ((1-v)I+vx^{-1})^{-1}, \end{aligned} \quad (20)$$

for any  $\nu \in [0, 1]$ .

If we multiply (6) both sides by  $A^{-1/2}$  we get  $MI \geq A^{-1/2}BA^{-1/2} \geq mI$ .

By writing the inequality (20) for  $X = A^{-1/2}BA^{-1/2}$  we obtain

$$\begin{aligned} & \left[ \nu(1-\nu) \left( 1 - \frac{\min\{M, 1\}}{\max\{m, 1\}} \right)^2 + 1 \right] \\ & \times \left( (1-\nu)I + \nu \left( A^{-1/2}BA^{-1/2} \right)^{-1} \right)^{-1} \\ & \leq (1-\nu)I + \nu A^{-1/2}BA^{-1/2} \\ & \leq \left[ \nu(1-\nu) \left( \frac{\max\{M, 1\}}{\min\{m, 1\}} - 1 \right)^2 + 1 \right] \\ & \times \left( (1-\nu)I + \nu \left( A^{-1/2}BA^{-1/2} \right)^{-1} \right)^{-1}, \end{aligned} \quad (21)$$

for any  $\nu \in [0, 1]$ .

If we multiply the inequality (21) both sides with  $A^{1/2}$ , then we get

$$\begin{aligned} & \left[ \nu(1-\nu) \left( 1 - \frac{\min\{M, 1\}}{\max\{m, 1\}} \right)^2 + 1 \right] \\ & \times A^{1/2} \left( (1-\nu)I + \nu \left( A^{-1/2}BA^{-1/2} \right)^{-1} \right)^{-1} A^{1/2} \\ & \leq (1-\nu)A + \nu B \\ & \leq \left[ \nu(1-\nu) \left( \frac{\max\{M, 1\}}{\min\{m, 1\}} - 1 \right)^2 + 1 \right] \\ & \times A^{1/2} \left( (1-\nu)I + \nu \left( A^{-1/2}BA^{-1/2} \right)^{-1} \right)^{-1} A^{1/2}, \end{aligned} \quad (22)$$

for any  $\nu \in [0, 1]$ .

Since

$$\begin{aligned} & A^{1/2} \left( (1-\nu)I + \nu \left( A^{-1/2}BA^{-1/2} \right)^{-1} \right)^{-1} A^{1/2} \\ & = A^{1/2} \left( (1-\nu)I + \nu A^{1/2}B^{-1}A^{1/2} \right)^{-1} A^{1/2} \\ & = A^{1/2} \left( A^{1/2} \left( (1-\nu)A^{-1} + \nu B^{-1} \right) A^{1/2} \right)^{-1} A^{1/2} \end{aligned}$$

$$\begin{aligned}
&= A^{1/2} \left( A^{-1/2} \left( (1-\nu) A^{-1} + \nu B^{-1} \right)^{-1} A^{-1/2} \right) A^{1/2} \\
&= \left( (1-\nu) A^{-1} + \nu B^{-1} \right)^{-1} = A!_\nu B
\end{aligned}$$

hence by (22) we get the desired result (16).  $\square$

We also have:

**Theorem 3** Let  $A, B$  be positive invertible operators and  $M > m > 0$  such that the condition (6) holds. Then we have

$$d_\nu(m, M) A!_\nu B \leq A \nabla_\nu B \leq D_\nu(m, M) A!_\nu B \quad (23)$$

for any  $\nu \in [0, 1]$ , where

$$d_\nu(m, M) := 4 \left[ \left( \nu - \frac{1}{2} \right)^2 + \nu(1-\nu) \times \begin{cases} K(M) & \text{if } M < 1, \\ 1 & \text{if } m \leq 1 \leq M, \\ K(m) & \text{if } 1 < m \end{cases} \right]$$

and

$$\begin{aligned}
D_\nu(m, M) &:= 4 \left[ \left( \nu - \frac{1}{2} \right)^2 + \nu(1-\nu) \times \begin{cases} K(m) & \text{if } M < 1, \\ \max\{K(m), K(M)\} & \text{if } m \leq 1 \leq M, \\ K(M) & \text{if } 1 < m. \end{cases} \right]
\end{aligned}$$

In particular, we have

$$d(m, M) A!B \leq A \nabla B \leq D(m, M) A!B \quad (24)$$

where

$$d(m, M) := \begin{cases} K(M) & \text{if } M < 1, \\ 1 & \text{if } m \leq 1 \leq M, \\ K(m) & \text{if } 1 < m \end{cases}$$

and

$$D(m, M) := \begin{cases} K(m) & \text{if } M < 1, \\ \max\{K(m), K(M)\} & \text{if } m \leq 1 \leq M, \\ K(M) & \text{if } 1 < m. \end{cases}$$

**Proof.** From (11) we have for any  $x \in (0, \infty)$  and for any  $\nu \in [0, 1]$  that

$$\frac{A_\nu(1, x)}{H_\nu(1, x)} - 1 = \nu(1-\nu) \frac{(x-1)^2}{x}. \quad (25)$$

Since  $K(x) - 1 = \frac{(x-1)^2}{4x}$ ,  $x > 0$ , then by (25) we have

$$\begin{aligned}\frac{A_\nu(1, x)}{H_\nu(1, x)} &= 1 + 4\nu(1-\nu)[K(x) - 1] \\ &= 4\nu(1-\nu)K(x) + 4\left(\nu - \frac{1}{2}\right)^2 \\ &= 4\left[\nu(1-\nu)K(x) + \left(\nu - \frac{1}{2}\right)^2\right]\end{aligned}$$

or, equivalently,

$$A_\nu(1, x) = 4\left[\nu(1-\nu)K(x) + \left(\nu - \frac{1}{2}\right)^2\right]H_\nu(1, x) \quad (26)$$

for any  $x \in (0, \infty)$  and for any  $\nu \in [0, 1]$ .

From (26) we then have for any  $x \in [m, M] \subset (0, \infty)$  that

$$\begin{aligned}4\left[\nu(1-\nu)\min_{x \in [m, M]} K(x) + \left(\nu - \frac{1}{2}\right)^2\right]H_\nu(1, x) \\ \leq A_\nu(1, x) \leq 4\left[\nu(1-\nu)\max_{x \in [m, M]} K(x) + \left(\nu - \frac{1}{2}\right)^2\right]H_\nu(1, x).\end{aligned} \quad (27)$$

Since

$$\min_{x \in [m, M]} K(x) = \begin{cases} K(M) & \text{if } M < 1, \\ 1 & \text{if } m \leq 1 \leq M, \\ K(m) & \text{if } 1 < m \end{cases}$$

and

$$\max_{x \in [m, M]} K(x) = \begin{cases} K(m) & \text{if } M < 1, \\ \max\{K(m), K(M)\} & \text{if } m \leq 1 \leq M, \\ K(M) & \text{if } 1 < m, \end{cases}$$

then by (27) we have

$$\begin{aligned}d_\nu(m, M) \left(1 - \nu + \nu x^{-1}\right)^{-1} &\leq 1 - \nu + \nu x \\ &\leq D_\nu(m, M) \left(1 - \nu + \nu x^{-1}\right)^{-1}\end{aligned} \quad (28)$$

for any  $x \in [m, M]$  and for any  $\nu \in [0, 1]$ .

By a similar argument to the one from Theorem 2 we deduce the desired operator inequality (23). The details are omitted.  $\square$

### 3 Some particular cases

Let  $A, B$  positive invertible operators and positive real numbers  $m, m', M, M'$  such that the condition  $0 < mI \leq A \leq m'I < M'I \leq B \leq MI$  holds.

Put  $h := \frac{M}{m}$  and  $h' := \frac{M'}{m'}$ , then we have

$$A < h'A = \frac{M'}{m'}A \leq B \leq \frac{M}{m}A = hA.$$

By (16) we get

$$\begin{aligned} \left[ v(1-v) \left( \frac{h'-1}{h'} \right)^2 + 1 \right] A!_v B &\leq A \nabla_v B \\ &\leq \left[ v(1-v)(h-1)^2 + 1 \right] A!_v B \end{aligned} \quad (29)$$

for any  $v \in [0, 1]$ .

By (23) we get

$$\begin{aligned} 4 \left[ \left( v - \frac{1}{2} \right)^2 + v(1-v) K(h') \right] A!_v B \\ \leq A \nabla_v B \leq 4 \left[ \left( v - \frac{1}{2} \right)^2 + v(1-v) K(h) \right] A!_v B \end{aligned} \quad (30)$$

for any  $v \in [0, 1]$ .

If  $0 < mI \leq B \leq m'I < M'I \leq A \leq MI$ , then for  $h := \frac{M}{m}$  and  $h' := \frac{M'}{m'}$  we also have

$$\frac{1}{h}A \leq B \leq \frac{1}{h'}A < A.$$

Finally, by (16) we get (29) while from (23) we get (30) as well.

## References

- [1] S. S. Dragomir, Bounds for the normalized Jensen functional, *Bull. Austral. Math. Soc.*, **74** (3) (2006), 417–478.
- [2] S. S. Dragomir, A note on Young's inequality, Preprint *RGMIA Res. Rep. Coll.*, **18** (2015), Art. 126. [Online <http://rgmia.org/papers/v18/v18a126.pdf>].

- 
- [3] S. S. Dragomir, Some new reverses of Young's operator inequality, Preprint *RGMIA Res. Rep. Coll.*, **18** (2015), Art. 130. [<http://rgmia.org/papers/v18/v18a130.pdf>].
  - [4] S. S. Dragomir, On new refinements and reverses of Young's operator inequality, Preprint *RGMIA Res. Rep. Coll.*, **18** (2015), Art. 135. [<http://rgmia.org/papers/v18/v18a135.pdf>].
  - [5] S. S. Dragomir, Some inequalities for operator weighted geometric mean, Preprint *RGMIA Res. Rep. Coll.*, **18** (2015), Art. 139. [<http://rgmia.org/papers/v18/v18a139.pdf>].
  - [6] S. S. Dragomir, Some reverses and a refinement of Hölder operator inequality, Preprint *RGMIA Res. Rep. Coll.* **18** (2015), Art. 147. [<http://rgmia.org/papers/v18/v18a147.pdf>].
  - [7] S. S. Dragomir, Upper and lower bounds for the difference between the weighted arithmetic and harmonic operator means, Preprint *RGMIA Res. Rep. Coll.*, **19** (2016), Art. [<http://rgmia.org/papers/v19/v19a0.pdf>].
  - [8] S. Furuichi, Refined Young inequalities with Specht's ratio, *J. Egyptian Math. Soc.*, **20** (2012), 46–49.
  - [9] S. Furuichi, On refined Young inequalities and reverse inequalities, *J. Math. Inequal.* **5** (2011), 21–31.
  - [10] W. Liao, J. Wu, J. Zhao, New versions of reverse Young and Heinz mean inequalities with the Kantorovich constant, *Taiwanese J. Math.*, **19** (2015), No. 2, pp. 467–479.
  - [11] M. Tominaga, Specht's ratio in the Young inequality, *Sci. Math. Japon.*, **55** (2002), 583–588.
  - [12] G. Zuo, G. Shi, M. Fujii, Refined Young inequality with Kantorovich constant, *J. Math. Inequal.*, **5** (2011), 551–556.

Received: October 11, 2018